

Plant Development



Edited by
Marja Timmerman



VOLUME NINETY ONE

CURRENT TOPICS IN
DEVELOPMENTAL BIOLOGY
Plant Development

Series Editor

Paul M. Wassarman
Department of Developmental and Regenerative Biology
Mount Sinai School of Medicine
New York, NY 10029-6574
USA

Olivier Pourquié
Institut de Génétique et de Biologie
Cellulaire et Moléculaire (IGBMC)
Inserm U964, CNRS (UMR 7104)
Université de Strasbourg
Illkirch
France

Editorial Board

Blanche Capel
Duke University Medical Center
Durham, NC, USA

B. Denis Duboule
Department of Zoology and Animal Biology
NCCR 'Frontiers in Genetics'
Geneva, Switzerland

Anne Ephrussi
European Molecular Biology Laboratory
Heidelberg, Germany

Janet Heasman
Cincinnati Children's Hospital Medical Center
Department of Pediatrics
Cincinnati, OH, USA

Julian Lewis
Vertebrate Development Laboratory
Cancer Research UK London Research Institute
London WC2A 3PX, UK

Yoshiki Sasai
Director of the Neurogenesis and Organogenesis Group
RIKEN Center for Developmental Biology
Chuo, Japan

Philippe Soriano
Department of Developmental and Regenerative Biology
Mount Sinai Medical School
New York, USA

Cliff Tabin
Harvard Medical School
Department of Genetics
Boston, MA, USA

Founding Editors

A. A. Moscona
Alberto Monroy

VOLUME NINETY ONE

CURRENT TOPICS IN
DEVELOPMENTAL BIOLOGY

Plant Development

Edited by

MARJA C. P. TIMMERMANS
Cold Spring Harbor Laboratory
Cold Spring Harbor
New York, USA



AMSTERDAM • BOSTON • HEIDELBERG • LONDON
NEW YORK • OXFORD • PARIS • SAN DIEGO
SAN FRANCISCO • SINGAPORE • SYDNEY • TOKYO
Academic Press is an imprint of Elsevier



Academic Press is an imprint of Elsevier
525 B Street, Suite 1900, San Diego, CA 92101-4495, USA
30 Corporate Drive, Suite 400, Burlington, MA 01803, USA
32, Jamestown Road, London NW1 7BY, UK
Linacre House, Jordan Hill, Oxford OX2 8DP, UK

First edition 2010

Copyright © 2010 Elsevier Inc. All rights reserved

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means electronic, mechanical, photocopying, recording or otherwise without the prior written permission of the publisher

Permissions may be sought directly from Elsevier's Science & Technology Rights Department in Oxford, UK: phone (+44) (0) 1865 843830; fax (+44) (0) 1865 853333; email: permissions@elsevier.com. Alternatively you can submit your request online by visiting the Elsevier web site at <http://elsevier.com/locate/permissions>, and selecting *Obtaining permission to use Elsevier material*

Notice

No responsibility is assumed by the publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the material herein. Because of rapid advances in the medical sciences, in particular, independent verification of diagnoses and drug dosages should be made

ISBN: 978-0-12-380910-0
ISSN: 0070-2153

For information on all Academic Press publications
visit our website at elsevierdirect.com

Printed and bound in the USA

10 11 12 13 9 8 7 6 5 4 3 2 1

Working together to grow
libraries in developing countries

www.elsevier.com | www.bookaid.org | www.sabre.org

ELSEVIER

BOOK AID
International

Sabre Foundation

CONTENTS

| | |
|--|-----------|
| <i>Contributors</i> | <i>xi</i> |
| <i>Preface</i> | <i>xv</i> |
| 1. Green Beginnings – Pattern Formation in the Early Plant Embryo | 1 |
| Cristina I. Llavata Peris, Eike H. Rademacher, and Dolf Weijers | |
| 1. Introduction–Seeds and Embryos | 2 |
| 2. Landmarks of Embryo Pattern Formation | 5 |
| 3. Regulation of Embryo Pattern Formation | 8 |
| 4. Evolutionary Aspects of Plant Embryogenesis | 20 |
| 5. Concluding Remarks | 22 |
| Acknowledgments | 22 |
| References | 23 |
| 2. Light-Regulated Plant Growth and Development | 29 |
| Chitose Kami, Séverine Lorrain, Patricia Hornitschek, and Christian Fankhauser | |
| 1. Multiple Photoreceptors to Sense a Variety of Light Colors and Intensities | 30 |
| 2. Physiological Responses Mediated by Plant Photoreceptors | 41 |
| 3. Photomorphogenesis in a Changing Environment | 47 |
| 4. Sites of Perception and Action of a Light Signal | 50 |
| 5. Signal Transduction | 52 |
| 6. Concluding Remarks | 58 |
| References | 58 |
| 3. Root Development–Two Meristems for the Price of One? | 67 |
| Tom Bennett and Ben Scheres | |
| 1. Introduction | 68 |
| 2. Structure of the Root | 68 |
| 3. Ontogeny of Roots | 70 |
| 4. The Root Stem Cell Niche | 76 |
| 5. The Meristematic Zone | 87 |
| 6. Elongation | 93 |
| | v |

| | |
|--|------------|
| 7. Differentiation | 95 |
| 8. Concluding Remarks | 96 |
| References | 96 |
| 4. Shoot Apical Meristem Form and function | 103 |
| Chan Man Ha, Ji Hyung Jun, and Jennifer C. Fletcher | |
| 1. Introduction | 104 |
| 2. Shoot Apical Meristem Organization | 104 |
| 3. Shoot Apical Meristem Maintenance | 107 |
| 4. Lateral Organ Initiation | 116 |
| 5. Phyllotaxis | 119 |
| 6. Communication Between the SAM and Differentiated Tissues | 122 |
| 7. Floral Meristem Termination | 124 |
| 8. Live Imaging and Computational Modeling Technology | 127 |
| 9. Concluding Remarks | 129 |
| References | 129 |
| 5. Signaling Sides: Adaxial–Abaxial Patterning in Leaves | 141 |
| Catherine A. Kidner and Marja C. P. Timmermans | |
| 1. Leaves Evolved Late in Land Plant History | 142 |
| 2. Leaves Develop from a Patterned Meristem | 142 |
| 3. Maintaining Polarity beyond the Meristem | 143 |
| 4. Determinants of Adaxial Fate— <i>PHANTASTICA</i> | 145 |
| 5. Determinants of Adaxial Fate—The HD-ZIPIII Genes | 147 |
| 6. Regulation of HD-ZIPIII Activity | 148 |
| 7. Determinants of Abaxial Fate—The KAN Genes | 150 |
| 8. Determinants of Abaxial Fate—The AUXIN RESPONSE FACTORS | 152 |
| 9. Regulation of <i>ARF3</i> and <i>ARF4</i> by the <i>TAS3</i> ta-siRNA Pathway | 152 |
| 10. How Are These Polarized Gene Expression Domains Established? | 155 |
| 11. Maintenance of Organ Polarity—Cross-Talk between Polarity Determinants | 157 |
| 12. Downstream Genes Regulating Medio-Lateral Blade Outgrowth | 158 |
| 13. Why So Complicated? | 160 |
| Acknowledgments | 162 |
| References | 162 |
| 6. Evolution Of Leaf Shape: A Pattern Emerges | 169 |
| Daniel Koenig and Neelima Sinha | |
| 1. Introduction | 170 |
| 2. Determinacy and Leaf Dissection | 170 |

| | |
|---|------------|
| 3. The Patterning of Growth During Leaf Development | 172 |
| 4. The Role of Auxin in Leaf Patterning | 174 |
| 5. What Distinguishes Lobes, Leaflets, and Blade? | 175 |
| 6. The Perception of Auxin Maxima and Specification of Differential Growth | 178 |
| 7. Homologies in Leaf Development | 179 |
| Acknowledgments | 181 |
| References | 181 |
| 7. Control of Tissue and Organ Growth in Plants | 185 |
| Holger Breuninger and Michael Lenhard | |
| 1. Introduction | 186 |
| 2. Description of Organ Growth | 187 |
| 3. Proliferative Growth Control | 190 |
| 4. Growth by Cell Elongation | 202 |
| 5. Compensation | 205 |
| 6. Coordination of Growth Across and Within Tissue Layers | 207 |
| 7. Modulation of Growth by Environmental Signals | 210 |
| 8. Modeling Plant Growth | 211 |
| 9. Concluding Remarks | 213 |
| References | 214 |
| 8. Vascular Pattern Formation in Plants | 221 |
| Enrico Scarpella and Ykä Helariutta | |
| 1. Introduction | 222 |
| 2. The Plant Vascular System | 223 |
| 3. Origin of Vascular Tissues | 225 |
| 4. Levels of Vascular Organization | 225 |
| 5. Control of Longitudinal Vascular Patterning and Integration with Organ Formation | 227 |
| 6. Regulation of Radial Vascular Patterning | 244 |
| 7. Concluding Remarks | 251 |
| Acknowledgments | 253 |
| Note Added in Proof | 253 |
| References | 253 |
| 9. Stomatal Patterning and Development | 267 |
| Juan Dong and Dominique C. Bergmann | |
| 1. Introduction | 268 |
| 2. The Signaling Cascade: From Extracellular Signals to Nuclear Factors | 270 |

| | |
|--|------------|
| 3. Signaling through a MAP Kinase Module | 278 |
| 4. Transcription Factors | 282 |
| 5. Regulation of Stomatal Asymmetric Cell Division | 288 |
| 6. Concluding Remarks | 293 |
| 7. Key Conclusions | 293 |
| References | 293 |
| | |
| 10. Trichome Patterning in <i>Arabidopsis thaliana</i>: From Genetic to Molecular Models | 299 |
| Rachappa Balkunde, Martina Pesch, and Martin Hülskamp | |
| 1. Introduction | 300 |
| 2. Trichome Form and Function in Plants | 300 |
| 3. Trichome Development in <i>Arabidopsis thaliana</i> | 301 |
| 4. Random Versus Regulated Trichome Pattern | 302 |
| 5. Genetic Analysis of the Core Patterning Machinery | 302 |
| 6. Genetic Interactions Between the Trichome Initiation Genes | 305 |
| 7. Expression Patterns of Patterning Genes: An Apparent Paradox | 306 |
| 8. Protein–Protein Interactions Between Patterning Components | 307 |
| 9. Transcriptional Cross-Regulation Between Patterning Genes | 308 |
| 10. Cellular Interactions are Mediated by Intercellular Protein Movement | 310 |
| 11. Immediate Downstream Genes—or Additional Patterning Components? | 311 |
| 12. The Role of Chromatin Structure in Patterning | 313 |
| 13. Modeling the Patterning System—Starting With Two Components | 314 |
| 14. Modeling the Patterning System: Global Interaction Models versus Solution of Discrete Problems | 316 |
| 15. Perspective | 317 |
| References | 317 |
| | |
| 11. Comparative Analysis of Flowering in Annual and Perennial Plants | 323 |
| Maria C. Albani and George Coupland | |
| 1. Introduction | 324 |
| 2. Control of Flowering in the Annual Model <i>Arabidopsis Thaliana</i> | 325 |
| 3. Flowering in Perennials | 333 |
| 4. The Evolution of Plant Life Strategies | 340 |
| 5. Concluding Remarks | 341 |
| References | 341 |

| | |
|--|------------|
| 12. Sculpting the Flower; the Role of microRNAs in Flower Development | 349 |
| Anwasha Nag and Thomas Jack | |
| 1. miRNAs in Plant Development | 350 |
| 2. miR172 Functions to Control Genes that Function in both Flowering Time Control and Floral Organ Identity in Arabidopsis | 354 |
| 3. miR172 Functions to Regulate Inflorescence Development in Maize | 361 |
| 4. miR156 Regulates Phase Change and Floral Induction | 363 |
| 5. miR164 Regulates Genes that Function in Organ Boundary Formation | 365 |
| 6. miR169 Represses C Class Activity in the Perianth in Antirrhinum and Petunia | 367 |
| 7. <i>miR319a</i> Controls Floral Organ Size and Shape in Arabidopsis | 368 |
| 8. miR159 Regulates Genes with Broad Roles in Flower Development | 369 |
| 9. miR167 is Critical for Male and Female Fertility | 370 |
| 10. Regulation of Floral Organ Polarity by miR166 | 371 |
| 11. Perspective and Future Challenges | 371 |
| Acknowledgments | 373 |
| References | 373 |
| | |
| 13. Development of Flowering Plant Gametophytes | 379 |
| Hong Ma and Venkatesan Sundaresan | |
| 1. Introduction | 380 |
| 2. Development of the Male Gametophyte or Pollen Grain | 380 |
| 3. Development of the Female Gametophyte or Embryo SAC | 391 |
| 4. Double Fertilization: Signaling and Reception between Male and Female | 400 |
| 5. Concluding Remarks | 403 |
| References | 404 |
| | |
| <i>Subject Index</i> | 413 |
| <i>Contents of Previous Volumes</i> | 423 |

This page intentionally left blank

CONTRIBUTORS

Maria C. Albani

Department of Plant Developmental Biology, Max Planck Institute for Plant Breeding Research, Cologne, Germany

Rachappa Balkunde

Botanical Institute III, University of Cologne, Köln, Germany

Tom Bennett

Department of Molecular Genetics, University of Utrecht, Utrecht, The Netherlands

Dominique C. Bergmann

Department of Biology, Stanford University, Stanford, California, USA

Holger Breuninger

Department of Cell & Developmental Biology, John Innes Centre, Norwich, UK

George Coupland

Department of Plant Developmental Biology, Max Planck Institute for Plant Breeding Research, Cologne, Germany

Juan Dong

Department of Biology, Stanford University, Stanford, California, USA

Christian Fankhauser

Center for Integrative Genomics, University of Lausanne, Lausanne, Switzerland

Jennifer C. Fletcher

Plant Gene Expression Center, USDA/UC Berkeley, Albany, California, USA and Department of Plant and Microbial Biology, University of California, Berkeley, California, USA

Chan Man Ha

Plant Gene Expression Center, USDA/UC Berkeley, Albany, California, USA and Department of Plant and Microbial Biology, University of California, Berkeley, California, USA

Ykä Helariutta

Department of Biological and Environmental Sciences, Institute of Biotechnology, University of Helsinki, Helsinki, Finland

Patricia Hornitschek

Center for Integrative Genomics, University of Lausanne, Lausanne, Switzerland

Martin Hülskamp

Botanical Institute III, University of Cologne, Köln, Germany

Thomas Jack

Department of Biological Sciences, Dartmouth College, Hanover,
New Hampshire, USA

Ji Hyung Jun

Plant Gene Expression Center, USDA/UC Berkeley, Albany, California,
USA and Department of Plant and Microbial Biology, University of California,
Berkeley, California, USA

Chitose Kami

Center for Integrative Genomics, University of Lausanne, Lausanne, Switzerland

Catherine A. Kidner

Royal Botanic Garden Edinburgh, Edinburgh, UK, and Institute of Molecular
Plant Sciences, University of Edinburgh, Edinburgh, UK

Daniel Koenig

Department of Plant Biology, University of California, Davis, California, USA

Michael Lenhard

Department of Cell & Developmental Biology, John Innes Centre, Norwich, UK

Cristina I. Llavata Peris

Laboratory of Biochemistry, Wageningen University, Wageningen,
The Netherlands

Séverine Lorrain

Center for Integrative Genomics, University of Lausanne, Lausanne, Switzerland

Hong Ma

State Key Laboratory of Genetic Engineering and School of Life Sciences, Institute
of Plant Biology, Fudan University, Shanghai, China, and Department of Biology,
Pennsylvania State University, University Park, Pennsylvania, USA

Anwasha Nag

Department of Biological Sciences, Dartmouth College, Hanover,
New Hampshire, USA

Martina Pesch

Botanical Institute III, University of Cologne, Köln, Germany

Eike H. Rademacher

Laboratory of Biochemistry, Wageningen University, Wageningen,
The Netherlands

Enrico Scarpella

Department of Biological Sciences, University of Alberta, Edmonton, Canada

Ben Scheres

Department of Molecular Genetics, University of Utrecht, Utrecht,
The Netherlands

Neelima Sinha

Department of Plant Biology, University of California, Davis, California, USA

Venkatesan Sundaresan

Departments of Plant Biology and Plant Sciences, University of California,
Davis, California, USA

Marja C. P. Timmermans

Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, USA

Dolf Weijers

Laboratory of Biochemistry, Wageningen University, Wageningen,
The Netherlands

This page intentionally left blank

PREFACE

This book captures many of the major advances made in the field of plant developmental biology. Over the past 20 years, genetic screens in *Arabidopsis* and other species have yielded a wide array of developmental mutants whose analyses have greatly enhanced our understanding of plant morphogenesis. Perhaps because of an initial focus on mutants with clearly defined, non-pleiotropic phenotypes, transcription factors were prominently represented among the first genes isolated from these screens. The field has matured to a point, however, that complex networks comprising signal transduction pathways, transcription factors, and downstream target genes are becoming clear for an ever-increasing number of developmental processes. Significant progress has been made on analysis of the mobile positional signals critical to plant development, which has long remained a main challenge in contemporary plant development. Moreover, by utilizing real-time analyses and computer modeling developmental principles are being elucidated in the context of the whole organ or organism.

Given the wealth of tools and resources available in *Arabidopsis*, the analysis of development has progressed most substantially in this model species. Comparative studies at the molecular and genetic level are beginning to illuminate the history of morphogenesis in plants, revealing evolutionary changes in key molecular players that form the basis for the enormous morphological diversity present within the plant kingdom. Moreover, with the tremendous advances in genomics, more complex systems are becoming accessible. This will allow new insights to be tested in the context of incisive classical experiments that have been the cornerstone of developmental biology, and can provide new interpretations of such formative experiments at the molecular and cellular level. Several of the reviews provide an insight into this new trend.

The chapters presented in this book describe our current knowledge, prospective insights, and key outstanding questions of multiple fundamental facets of plant development. Chapter 1 outlines the formative events that sculpt the organization of the plant body during embryogenesis and discusses the molecular mechanisms that regulate these patterning processes. It emerges that auxin has a profound coordinating role in embryogenesis. One example of this is in the formation of the embryonic root (Chapter 3), where an auxin reflux loop specifies the stem cell niche. This chapter also presents the progress in our understanding of root meristem function, outlining the distinctive properties of stem cells within the root meristem

and the network that controls root growth through the coordinate regulation of cell division and elongation. Germination as well as numerous other developmental transitions and adaptive growth responses are regulated through the perception of light. How plants sense light and coordinate the molecular events underlying photomorphogenesis is presented in Chapter 2. The signaling networks that maintain the position and activity of stem cells within the shoot apical meristem and that organize the initiation of lateral organs on the meristem periphery are discussed in Chapter 4. The complexity of these networks is remarkable, but with the development of live imaging approaches and computational tools, our understanding of the central principles of shoot apical meristem function has rapidly advanced.

The next three chapters focus on patterning, growth, and the morphological diversification of lateral organs in plants. Chapter 5 describes the evolutionary distinct contributions of three highly conserved interlocking pathways to leaf polarity and outlines the unique contributions of mobile small RNAs as positional signals that refine the boundary between the upper and lower sides of the leaf. Chapter 6 compares the mechanisms that regulate leaf shape in simple and compound leafed model species. Auxin is recognized as a key player also in this aspect of development. The emergent picture is one in which developmental pathways that modulate determinacy within the leaf can vary auxin-signaling outputs to generate diverse compound and simple leaf forms. Recent insights into how plant organs reach their characteristic final shape and size are presented in Chapter 7. In addition to the genetic networks that affect organ size at the cellular level by regulating cell proliferation and expansion, insights into the systems that coordinate such cellular processes across the entire organ or modulate them in response to environmental stimuli are discussed.

New insights into the patterning processes that give rise to the precise arrangements of vasculature, stomatal complexes, and trichomes are outlined in Chapters 8 through 10. Interestingly, these tissue or cell patterns are generated through entirely distinct mechanisms. Chapter 8 describes the role of polarized auxin transport in generating the distinctive arrangements of vascular strands within leaves, stems, and roots. In contrast, stereotyped asymmetric cell divisions are at the center of stomatal patterning and, as outlined in Chapter 9, the spacing of such instructive divisions is regulated by signaling pathways involving secreted peptide ligands and receptor-like kinases. Finally, Chapter 10 discusses current models for trichome patterning, presenting evidence for the involvement of two parallel spacing mechanisms; a lateral inhibition mechanism that relies in part on plasmodesmatal movement of trichome inhibitor proteins and an activator-depletion mechanism in which inducers of trichome differentiation are depleted around incipient trichome cells.

Chapter 11 reviews our understanding of the networks that regulate the switch from vegetative growth to flowering in response to intrinsic signals and environmental stimuli in *Arabidopsis thaliana*, and then broadens the discussion to the control of flowering in perennial species. The contribution of miRNAs in regulating the transition to flowering is described in Chapter 12. This chapter also outlines the varied contributions of small regulatory RNAs to the specification of floral organ identity and the control of floral organ shape and size. The final chapter of the book focuses on the progress in understanding the development of the male and the female gametophytes, and their interactions during the process of double fertilization. Signaling molecules and downstream genetic pathways that direct the differentiation of distinct cell types within the developing gametophytes, pollen tube attraction, or fertilization are beginning to emerge. An exciting example is the recent finding that auxin acts as a morphogenic determinant in the patterning of the female gametophyte.

Besides providing timely reviews of new advances in many fundamental aspects of plant development, the chapters presented in this book illustrate clearly the diversity of patterning mechanisms that drive plant morphogenesis. Mobile transcription factors, secreted peptides, hormones, and small regulatory RNAs function as instructive signals feeding into a wide array of signaling pathways and gene regulatory networks. This wealth of new data also allows us to draw parallels with animal development. It is now probably safe to state, rather than speculate, that many of the individual molecular components are either different or wired differently, but that the logic of developmental circuitry is conceptually quite similar between plants and animals. This book is testimony to the progress in our understanding of plant development that has been made in a surprisingly short time.

This page intentionally left blank

GREEN BEGINNINGS – PATTERN FORMATION IN THE EARLY PLANT EMBRYO

Cristina I. Llavata Peris,* Eike H. Rademacher,* and Dolf Weijers

Contents

| | |
|---|----|
| 1. Introduction—Seeds and Embryos | 2 |
| 2. Landmarks of Embryo Pattern Formation | 5 |
| 2.1. Formation of the apico-basal axis | 5 |
| 2.2. Outside versus inside—radial pattern establishment | 7 |
| 2.3. Stem cell niches—sources of all other cells | 7 |
| 2.4. Initiation of cotyledons | 7 |
| 3. Regulation of Embryo Pattern Formation | 8 |
| 3.1. Auxin biosynthesis, transport, and response | 8 |
| 3.2. Specification of the apico-basal axis | 10 |
| 3.3. Establishment of the radial pattern | 13 |
| 3.4. Initiation of meristems | 16 |
| 3.5. Cotyledon initiation | 18 |
| 4. Evolutionary Aspects of Plant Embryogenesis | 20 |
| 5. Concluding Remarks | 22 |
| Acknowledgments | 22 |
| References | 23 |

Abstract

Embryogenesis in plants transforms the zygote into a relatively simple structure, the seedling, which contains all tissues and organs that later form the mature plant body. Despite a profound diversity in cell division patterns among plant species, embryogenesis yields remarkably homologous seedling architectures. In this review, we describe the formative events during plant embryogenesis and discuss the molecular mechanisms that regulate these processes, focusing on *Arabidopsis*. Even though only a relatively small number of factors are known that regulate each patterning step, a picture emerges where locally acting transcription factors and intercellular signaling contribute to the specification and spatio-temporal coordination of the various cell types in the embryo.

Laboratory of Biochemistry, Wageningen University, Wageningen, The Netherlands

* These authors have contributed equally to this manuscript.

Current Topics in Developmental Biology, Volume 91
ISSN 0070-2153, DOI 10.1016/S0070-2153(10)91001-6

© 2010 Elsevier Inc.
All rights reserved.

Notably, several patterning processes are controlled by the plant hormone auxin. Most regulators that were identified in *Arabidopsis* have orthologs in other sequenced plant genomes, and several of these are expressed in similar patterns. Therefore, it appears that robust conserved mechanisms may underlie pattern formation in plant embryos.

1. INTRODUCTION—SEEDS AND EMBRYOS

During evolution, plants as largely immobile organisms have developed sophisticated mechanisms to disperse their offspring and explore new habitats. One way to expand the populated area is to grow into the surrounding by means of vegetative propagation via specialized structures such as rhizomes or stolons. However, these mechanisms do not reach over great distances and fail to overcome relatively small obstacles.

The innovation that surpassed such restrictions and allowed plants to use wind, water, animals, gravity, and even ballistic mechanisms as means of dispersal is the seed. Furthermore, seeds permit the enclosed embryo to remain in a dormant state and survive long periods of harsh environmental conditions, thereby mastering not only spatial but also temporal restrictions to its habitat. Finally, when conditions are beneficial for germination, the seed provides the embryo with the nutrients that support the first steps of post-embryonic development. To fulfill all of these different requirements, seeds develop in very different shapes and sizes. Adaptations in seed morphology have enabled plants to occupy new ecological niches with the result that plants now dominate habitats on all different kinds of land, in both hot and cold climates (Kessler and Stuppy, 2006).

Seed-bearing plants (spermatophytes) can be subdivided into two major groups: gymnosperms and angiosperms. The name-giving difference between these lies in the organization of the reproductive organs. Gymnosperms bear ovules on scales, which are usually in cone-like structures as, for example, in pine. Angiosperms instead have their reproductive organs arranged in flowers. Another important difference is that flowering plants feature a double fertilization step to produce the (typically) diploid embryo and the triploid extra-embryonic endosperm (Lersten, 2004). Despite these differences in seed anatomy and function, the development and basic body organization of the embryo are very similar among most higher plants (Cairney and Pullman, 2007; Johri *et al.*, 1992). Accordingly, embryogenesis in both groups undergoes three common phases. Initially the fertilized egg cell (zygote) undergoes elongation and after a few cell division rounds develops an apico-basal and radial axis. During the next phase, further cell divisions occur and the primordia of the fundamental organs for post-embryonic growth are established. A final phase of desiccation prepares the mature

embryo for dormancy and equips the seed with storage products that aid the seedling in the process of germination (Bewley and Black, 1994).

Typically, seeds carry a single embryo that emerges as the product of fertilization of the egg cell (Fig. 1.1, box 1). However, the formation of poly-embryonic seeds has been observed in many taxa (e.g., *Rutacea* (Citrus) or *Pinacea* (Pinus); Batygina and Vinogradova, 2007; Lakshmanan and Ambegaokar, 1984). Interestingly, the additional embryos within seeds do not always originate from cells that are derived from the zygote. More importantly, this is not even obligatory in mono-embryonic seeds. Hence, also unfertilized cells possess the potential to form embryos within a seed. Such asexual reproductive mechanisms of embryo formation have been summarized under the term *apomixis* (Fig. 1.1, box 2). We refer the reader to another review (Koltunow and Grossniklaus, 2003) that covers the various developmental origins of apomictic embryos. In addition to the naturally occurring origins of embryogenesis, male gametes (pollen grains) can give rise to viable but haploid embryos when cultured under certain




Figure 1.1 Origins of plant embryos. Embryogenesis normally occurs in the reproductive tissues within the flower. Shown here is a flower with subtending leaf. Zygotic embryogenesis (box 1) occurs when a pollen grain lands on the stigma of the carpel and fertilizes the egg cell in an ovule contained within the gynoecium. This ovule develops into a seed (brown oval) which, upon germination, generates the seedling (here dicotyledonous). Alternative modes of embryogenesis include the initiation of apomictic embryos within the ovule (box 2), microspore-derived embryos developing from pollen grains (box 3), and somatic embryos that are initiated from callus tissue derived from somatic cells (box 4). While only zygotic and apomictic embryogenesis pass through a seed stage, all these different modes of embryogenesis finally give rise to seedlings with the same body plan.




Figure 1.2 Fertilization in *Arabidopsis*. Fertilization occurs within the gynoecium (A), where ovules (B) are attached to the placental tissues. The *Arabidopsis* ovule consists of a gametophyte (embryo sac) covered with somatic integument cell layers. The egg cell, surrounded by two synergid cells, is localized at the micropylar end. Additionally, the embryo sac houses a central cell nucleus and three antipodal cells. (C) Upon germination and growth, the pollen tube is attracted to the ovule by the synergid cells. The pollen tube then releases its generative nuclei in the degenerating synergid, upon which the nuclei fuse with the egg cell and central cell nuclei. (B, C) Drawn after Sundaresan and Alandete-Saez (2010).

conditions (Seguí-Simarro and Nuez, 2008; Fig. 1.1, box 3). Finally, somatic cells can be forced into an embryogenic pathway by treatment of cultured explants with auxin (2,4-D) (Toonen and de Vries, 1996; Fig. 1.1, box 4).

As transpires from the above, a wide variety of cells can generate embryos either naturally or after experimental treatment. However, despite the distinct origins of embryos and the often different patterns of cell divisions within the embryo, the outcome is always a seedling that consists of an apico-basal axis with apical shoot and root meristems, a radial axis that defines the respective tissues, and one or two cotyledons positioned in the immediate surrounding of the shoot apical meristem (SAM). It is therefore likely that a robust set of conserved genetically encoded instructions underlies pattern formation during embryogenesis in all species.

Molecular processes underlying pattern formation and development are best described in *Arabidopsis thaliana*. Due to the regular pattern of cell divisions during embryogenesis (Jürgens and Mayer, 1994), this species has also become the model for studying zygotic embryogenesis. In this review we will discuss embryogenesis in *Arabidopsis* to exemplify basic mechanisms that sculpt the organization of the plant body during embryogenesis. In the following sections, we will first describe the developmental landmarks during embryogenesis, and then discuss the molecular mechanisms that control pattern formation, followed by a perspective on the evolutionary conservation of the regulatory mechanisms that have been found in *Arabidopsis*.

2. LANDMARKS OF EMBRYO PATTERN FORMATION

In angiosperms such as *Arabidopsis*, the female gamete (egg cell) is positioned within the embryo sac, which in turn is embedded in the protective maternal tissue of the ovule inside the carpel (Fig. 1.2A, B). After deposition of a male microgametophyte (pollen grain) on the stigma of the carpel, a pollen tube extends toward the ovule. This tube eventually enters the ovule via its micropylar end and facilitates the delivery of two haploid sperm nuclei. One of these nuclei fertilizes the egg cell while the other fuses with the two nuclei of the central cell to give rise to the triploid endosperm (Lersten, 2004; Fig. 1.2C). Embryo and endosperm develop simultaneously, and several mutant studies suggest that their growth patterns are interdependent (reviewed in Berger *et al.*, 2006). So far, however, there is no conclusive evidence that embryo pattern formation depends on the presence of the endosperm. Hence, consistent with the flexible origin of embryos, pattern formation in the embryo is driven by intrinsic factors rather than by environmental cues.

Embryo patterning follows a series of “landmarks” that each establishes part of the final organization. In the following, we briefly describe each such landmark.

2.1. Formation of the apico-basal axis

Embryo development occurs within the highly polarized environment of the ovule. Both the ovule and the embryo sac have a distinct polarized axis, and also the egg cell is intrinsically polar as judged from the localization of its organelles on the basal (e.g. vacuole) or apical end (e.g. nucleus) (Fig. 1.2C; Lersten, 2004). Fertilization induces stretching of the zygote, which is followed by an asymmetric division that gives rise to two daughter cells with different composition, shape, and developmental fate. The apical cell is small with a dense cytoplasm, while the basal cell is large and vacuolated. With three rounds of cell divisions, the apical cell generates a spherical proembryo consisting of a total of eight cells, while the basal cell only divides transversally and gives rise to a transient filamentous structure called the suspensor. This extra-embryonic suspensor connects the proembryo to maternal tissue and pushes it into the lumen of the ovule. Later during embryogenesis, only the uppermost suspensor cell, the hypophysis, becomes incorporated in the embryonic root meristem, as precursor of the quiescent center (QC) and central root cap cells (see below) (Fig. 1.3A).

At the eight-cell (octant) stage, four different domains can be distinguished along the apico-basal axis of the embryo; the proembryo consists of an upper and a lower tier of four cells each and is positioned on top of hypophysis and suspensor (Fig. 1.3A, B).

- [read Balzac journaliste : Articles et chroniques pdf, azw \(kindle\)](#)
- **[Child Soldier book](#)**
- **[read online Buddhism in Sinhalese Society 1750-1900: A Study of Religious Revival and Change](#)**
- [download online *The Wisdom of the Enneagram: The Complete Guide to Psychological and Spiritual Growth for the Nine Personality Types*](#)

- <http://monkeybubblemedia.com/lib/Robert-Ludlum-s-The-Lazarus-Vendetta--Covert-One--Book-5-.pdf>
- <http://rodrigocaporal.com/library/Every-Man-for-Himself.pdf>
- <http://pittiger.com/lib/Mandela-s-Way--Fifteen-Lessons-on-Life--Love--and-Courage.pdf>
- <http://dadhoc.com/lib/Philosophy--A-Beginner-s-Guide.pdf>