

Seshadev Padhi  
Smita Pati

# Theory of Third-Order Differential Equations

 Springer

---

# Theory of Third-Order Differential Equations

---

Seshadev Padhi • Smita Pati

# Theory of Third-Order Differential Equations

 Springer

---

Seshadev Padhi  
Department of Applied Mathematics  
Birla Institute of Technology, Mesra  
Ranchi, Jharkhand, India

Smita Pati  
Department of Applied Mathematics  
Birla Institute of Technology, Mesra  
Ranchi, Jharkhand, India

ISBN 978-81-322-1613-1                      ISBN 978-81-322-1614-8 (eBook)  
DOI 10.1007/978-81-322-1614-8  
Springer New Delhi Heidelberg New York Dordrecht London

Library of Congress Control Number: 2013951151

© Springer India 2014

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed. Exempted from this legal reservation are brief excerpts in connection with reviews or scholarly analysis or material supplied specifically for the purpose of being entered and executed on a computer system, for exclusive use by the purchaser of the work. Duplication of this publication or parts thereof is permitted only under the provisions of the Copyright Law of the Publisher's location, in its current version, and permission for use must always be obtained from Springer. Permissions for use may be obtained through RightsLink at the Copyright Clearance Center. Violations are liable to prosecution under the respective Copyright Law.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

While the advice and information in this book are believed to be true and accurate at the date of publication, neither the authors nor the editors nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Printed on acid-free paper

Springer is part of Springer Science+Business Media ([www.springer.com](http://www.springer.com))

---

*Dedicated to Suchitra and Aryadev*

---

# Preface

Over the last four decades, intensive work has been carried out in the field of theory of nonautonomous third-order ordinary and delay differential equations. However, it is only recently that the global attractivity of third-order equations has been given a serious study. During these years, new investigations were developed and results of principal importance were obtained. In particular, suitable oscillation criteria for third-order linear and nonlinear differential equations were established, with emphasis on the oscillation and nonoscillation of third-order homogeneous differential equations. Existence and nonexistence conditions for oscillatory and nonoscillatory solutions of various types were found and asymptotic formulae for solutions of a sufficiently wide class of linear and nonlinear equations were derived. These studies notwithstanding, one would observe that oscillation and asymptotic behaviour of nonoscillatory solutions of third-order delay differential equations has rarely been studied.

In this book, an attempt is made to sum up these results. The necessity of such an attempt is felt especially as the well-known monograph of M. Greguš, *Third Order Linear Differential Equations*, Reidel, Dordrecht, The Netherlands, 1982, devoted to related topics, reflects results which are 30 years old.

This book consists of seven chapters. In the first one, we consider the third-order linear differential equation, with constant coefficients, of the form

$$x''' + ax'' + bx' + cx = 0, \quad (1)$$

where  $a, b$  and  $c$  are constants. Eight different cases of  $a, b$  and  $c$  were considered while dealing with (1). Further, it is observed that different structures of solution spaces of (1) appears for different cases on  $a, b$  and  $c$ . In this chapter, an introduction is given to the oscillation theory of the nonhomogeneous equation

$$x''' + ax'' + bx' + cx = f, \quad (2)$$

where  $a, b, c$  and  $f$  are constants. An apparatus of comparison theorems is developed, which allows us to establish criteria for (2) to have oscillatory or nonoscillatory solutions with the help of oscillation and nonoscillation of (1). An overall

idea on oscillation of third-order delay differential equations is also included in this chapter. Additionally, some basic results are incorporated, which are needed in the forthcoming chapters.

Chapter 2 deals with the linear equation

$$x''' + a(t)x'' + b(t)x' + c(t)x = 0, \quad (3)$$

where  $a, b \in C^1([\sigma, \infty), R)$  and  $c \in C([\sigma, \infty), R)$ ,  $\sigma \in R$ . This chapter contains seven sections, where most of the results obtained for (1) have been generalised to (3). Many problems have been proposed in the sections or at the end of the chapter. Apart from these, some new nontrivial sufficient conditions for the oscillation of (3) are given. Further, several sufficient conditions, different from earlier ones, are given which can be applied to the third-order Euler equation

$$x''' + \frac{a_0}{t}x'' + \frac{b_0}{t^2}x' + \frac{c_0}{t^3}x = 0,$$

where  $a_0, b_0$  and  $c_0$  are real constants. Several comparison principles have been used to establish the nonoscillation of (3). We have established criteria for (3) to have a family of solutions asymptotically, equivalent to solutions of (3), and we studied the properties of this family. Finally, existence criteria for solutions vanishing at infinity have been explored in this chapter.

Chapter 3 is concerned with the oscillation of the equation

$$x''' + a(t)x'' + b(t)x' + c(t)x = f(t), \quad (4)$$

where  $a(t), b(t)$  and  $c(t)$  are as defined earlier and  $f \in C([\sigma, \infty), R)$ . Some of the eight different cases of  $a(t), b(t)$  and  $c(t)$  have been considered while the remaining have been retained as open problems. We have applied the results of Chap. 2 to obtain sufficient conditions for the oscillation and nonoscillation of (4).

In Chap. 4, emphasis is given to the oscillation and nonoscillation of solutions and the asymptotic behaviour of the third-order nonlinear differential equations

$$\begin{aligned} x''' + b(t)x' + c(t)x^\alpha &= 0, \\ x''' + b(t)x' + c(t)f(x) &= 0, \end{aligned}$$

and results of more general equations

$$x''' + a(t)x'' + b(t)x' + c(t)x^\alpha = 0,$$

and

$$x''' + a(t)x'' + b(t)x' + c(t)f(x) = 0,$$

where  $a(t), b(t)$  and  $c(t)$  were defined earlier in (4) and  $f : R \rightarrow R$  and some other restrictions on  $f$ . In this chapter, we solve the Kneser problem on monotone solutions and prove, under certain restrictions on  $f$  and  $\alpha$ , that monotone solutions tend to zero eventually.

Chapter 5 is quite interesting. We have studied oscillation of solutions of the third-order nonlinear and nonhomogeneous equations of the form

$$(r(t)x'')' + q(t)(x')^\beta + p(t)x^\alpha = f(t), \quad (5)$$

where  $r, p, q$  and  $f \in C([\sigma, \infty), R)$  and  $r(t) > 0$ ,  $\alpha$  and  $\beta$  are quotient of odd integers. Along the way, we have given some interesting results on oscillation of the nonlinear equation

$$x''' + a(t)x'' + b(t)x' + c(t)x = f(t, x, x', x''), \quad (6)$$

where  $a, b$  and  $c \in C([\sigma, \infty), R)$  and  $f : [\sigma, \infty) \times R^3 \rightarrow R$ .

In Chap. 6, we apply the results of Chap. 2 to obtain oscillation and asymptotic behaviour of nonoscillation of the third-order delay differential equation

$$x'''(t) + a(t)x''(t) + b(t)x'(t) + c(t)x(g(t)) = 0, \quad (7)$$

where  $a(t), b(t)$  and  $c(t)$  are as defined earlier and  $g : R \rightarrow R$  with the property that  $0 \leq g(t) \leq t$  and  $\lim_{t \rightarrow \infty} g(t) = \infty$ . It has been observed that the presence of the delay term  $g(t)$  makes it difficult to make a direct study of the asymptotic behaviour of solutions of (7). However, the canonical transformations makes it possible to study the properties of solutions of (7). Equations (3) and (7) have been transformed to equivalent canonical equations and then the asymptotic behaviour of solutions of (7) have been studied by knowing the oscillation and nonoscillation of (3). We have classified the nonoscillatory solutions of (7) as Property A and Property B, according to their asymptotic behaviour.

In the last chapter, we study the global attractivity of solutions of equations

$$x''' + \psi(x, x')x'' + f(x, x') = 0 \quad (8)$$

and

$$x''' + \psi(x, x')x'' + f(x, x') = p(t), \quad (9)$$

where  $\psi, f, \psi_x, f_x \in C(R \times R, R)$  and  $p \in C([0, \infty), R)$ . Lyapunov function has been used to obtain the results. A direct step by step method has been used to find the asymptotic stability of solutions of the third-order equation

$$\begin{aligned} x'''(t) = & p_1x''(t) + p_2x''(t - \tau) + q_1x'(t) + q_2x'(t - \tau) + r_1x(t) \\ & + r_2x(t - \tau), \quad t \geq 0, \end{aligned} \quad (10)$$

with the initial condition

$$x(t) = \phi(t), \quad t \in [-\tau, 0], \quad (11)$$

where  $p_1, p_2, q_1, q_2, r_1$  and  $r_2$  are real constants,  $\tau > 0$  is a real number and  $\phi \in C([-\tau, 0), R)$  is an initial function. The results are then applied to obtain stability and asymptotic stability of solutions of (11).

Some results are given in the form of open problems to be solved by the readers. We also state some problems whose solutions we do not know.

Seshadev Padhi  
Smita Pati



---

## Acknowledgements

It is almost impossible to thank everyone who helped us over the years to understand the theory of third-order differential equations. However, three people must be singled out: Prof. N. Parhi who taught us oscillation theory of third-order differential equations, Prof. John R. Graef for his constant inspiration and valuable suggestions, and Prof. Chuanxi Qian who inspired us to learn stability theory of third-order differential equations. We are also thankful to Prof. Julio G. Dix, Prof. E. Thandapani, Prof. P.D.N. Srinivasu and Dr. B.S.R.V. Prasad for their constant inspiration and encouragement in preparation of this book. We should also mention here that it was a great pleasure to work with Mr. Shamim Ahmad and Ms. Nupoor Singh, the editorial team of Springer and the production team, especially Mr. Donatas Akmanavičius, who took utmost care during the preparation of the book. We are thankful to our family members, especially to Suchitra Panigrahi, Aryadev Padhi, Mr. Bansidhar Padhi, Mr. Sitanath Panigrahi and Mr. Ashok Kumar Pati, for their constant inspirations in preparing this book.

---

# Contents

<b>1</b>	<b>Introduction</b> . . . . .	<b>1</b>
1.1	Preliminaries . . . . .	1
1.2	Third-Order Linear Differential Equations with Constant Coefficients . . . . .	8
1.3	Introduction to Third-Order Delay Differential Equations . . . . .	19
1.4	Introduction to Third-Order Canonical Differential Equations . . . . .	22
1.5	Some Basic Results . . . . .	29
1.6	Some Useful Results from Analysis . . . . .	39
1.7	Notes . . . . .	41
	References . . . . .	41
<b>2</b>	<b>Behaviour of Solutions of Linear Homogeneous Differential Equations of Third Order</b> . . . . .	<b>45</b>
2.1	Behaviour of Solutions of $x''' + a(t)x'' + b(t)x' + c(t)x = 0$ with $a(t) \geq 0, b(t) \leq 0$ and $c(t) > 0$ . . . . .	46
2.2	Behaviour of Solutions of $x''' + a(t)x'' + b(t)x' + c(t)x = 0$ with $a(t) \leq 0, b(t) \leq 0$ and $c(t) > 0$ . . . . .	65
2.3	Behaviour of Solutions of $x''' + a(t)x'' + b(t)x' + c(t)x = 0$ with $a(t) \leq 0, b(t) \leq 0$ and $c(t) < 0$ . . . . .	73
2.4	Behaviour of Solutions of $x''' + a(t)x'' + b(t)x' + c(t)x = 0$ with $a(t) \geq 0, b(t) \leq 0$ and $c(t) < 0$ . . . . .	83
2.5	Behaviour of Solutions of $x''' + a(t)x'' + b(t)x' + c(t)x = 0$ with $a(t) \geq 0, b(t) \geq 0$ and $c(t) > 0$ . . . . .	93
2.6	Behaviour of Solutions of $x''' + a(t)x'' + b(t)x' + c(t)x = 0$ with $a(t) \leq 0, b(t) \geq 0$ and $c(t) < 0$ . . . . .	107
2.7	Oscillation and Nonoscillation of Third-Order Linear Differential Equations of the Form $(r_2(t)(r_1(t)x')')' + r_3(t)x = 0$ . . . . .	119
2.8	Open Problems and Discussions . . . . .	143
2.9	Notes . . . . .	145
	References . . . . .	145

<b>3</b>	<b>Oscillation of Solutions of Linear Nonhomogeneous Differential Equations of Third Order</b>	<b>147</b>
3.1	Nonoscillatory Behaviour of Solutions of Nonhomogeneous Linear Differential Equations	148
3.2	Oscillatory Behaviour of Solutions of Nonhomogeneous Linear Differential Equations	169
3.3	Nonoscillation of Nonhomogeneous Linear Differential Equations	177
3.4	Asymptotic Behaviour of Oscillatory Solutions of Third-Order Linear Nonhomogeneous Differential Equations	184
3.5	Open Problems and Discussions	186
3.6	Notes	190
	References	190
<b>4</b>	<b>Oscillation and Nonoscillation of Homogeneous Third-Order Nonlinear Differential Equations</b>	<b>193</b>
4.1	Behaviour of Solutions of $x''' + b(t)x' + c(t)x^\alpha = 0$ and $x''' + b(t)x' + c(t)f(x) = 0$	197
4.2	Behaviour of Solutions of $x''' + a(t)x'' + b(t)x' + c(t)x^\alpha = 0$ and $x''' + a(t)x'' + b(t)x' + c(t)f(x) = 0$	239
4.3	Behaviour of Solutions of $x''' + c(t)x^\alpha = 0$ and $x''' + c(t)f(x) = 0$	250
4.4	Behaviour of Solutions of $(r_2(t)(r_1(t)x')')' + q(t)x' + p(t)x^\alpha = 0$ and $(r_2(t)(r_1(t)x')')' + q(t)x' + p(t)f(x) = 0$	277
4.5	Open Problems and Discussions	300
4.6	Notes	301
	References	301
<b>5</b>	<b>Oscillation and Nonoscillation of Nonlinear Nonhomogeneous Differential Equations of Third Order</b>	<b>305</b>
5.1	Oscillatory and Nonoscillatory Behaviour of Solutions of $x''' + a(t)x'' + b(t)x' + c(t)x = f(t, x, x', x'')$	306
5.2	Nonoscillation of Solutions of $(r_2(t)(r_1(t)x')')' + q(t)(x')^\beta + p(t)x^\alpha = f(t)$	319
5.3	Open Problems and Discussions	334
5.4	Notes	334
	References	334
<b>6</b>	<b>Oscillatory and Asymptotic Behaviour of Solutions of Third-Order Delay Differential Equations</b>	<b>335</b>
6.1	Asymptotic Behaviour of Solutions of Linear Delay Differential Equations of the Form $x'''(t) + a(t)x''(t) + b(t)x'(t) + c(t)x(g(t)) = 0$	336
6.2	Oscillation and Asymptotic Properties of Solutions of the Linear Delay Differential Equation $\frac{1}{r_3(t)}(\frac{1}{r_2(t)}(\frac{1}{r_1(t)}(\frac{x(t)}{r_0(t)})')')' + p(t)x(g(t)) = 0$	378

6.3	Oscillation and Asymptotic Behaviour of Solutions of the Linear Delay Differential Equation $x'''(t) + p(t)x(g(t)) = 0$ . . . . .	400
6.4	Asymptotic Behaviour of Solutions of Nonlinear Delay-Differential Equation of the Form $x'''(t) + q(t)x'(t) + p(t)f(x(g(t))) = 0$ . . . . .	414
6.5	Oscillation of Solutions of the Nonlinear Delay Differential Equation $(r(t)(x''(t))^\gamma)' + p(t)x^\gamma(g(t)) = 0$ and $(r(t)(x''(t))^\gamma)' + p(t)f(x(g(t))) = 0$ . . . . .	421
6.6	Oscillation and Asymptotic Behaviour of Solutions of $(r_2(t)(r_1(t)x'(t))^\gamma)' + q(t)x'(t) + p(t)f(x(g(t))) = 0$ . . . . .	428
6.7	Oscillation of Solutions of Third-Order Differential Equations with Distributed Deviating Arguments . . . . .	435
6.8	Oscillation and Asymptotic Behaviour of Nonoscillatory Solutions of $(r_2(t)(r_1(t)x'(t))^\gamma)' + p(t)f(x(g(t))) = 0$ . . . . .	445
6.9	Nonoscillation of Solutions of Nonlinear Nonhomogeneous Delay Differential Equations of Third Order . . . . .	448
6.10	Open Problems and Discussions . . . . .	450
6.11	Notes . . . . .	451
	References . . . . .	451
<b>7</b>	<b>Stability of Third-Order Differential Equations</b> . . . . .	<b>455</b>
7.1	Stability of Solutions of $x''' + \psi(x, x')x'' + f(x, x') = 0$ . . . . .	456
7.2	Stability of Solutions of $x''' + \psi(x, x')x'' + f(x, x') = p(t)$ . . . . .	465
7.3	Stability of Solutions of $x''' + \psi(x, x', x'')x'' + f(x, x') = p(t, x, x', x'')$ . . . . .	474
7.4	Stability of Solutions of $x'''(t) = p_1x''(t) + p_2x''(t - \tau) + q_1x'(t) + q_2x'(t - \tau) + r_1x(t) + r_2x(t - \tau)$ . . . . .	481
7.5	Open Problems and Discussions . . . . .	501
7.6	Notes . . . . .	502
	References . . . . .	502
	<b>Index</b> . . . . .	<b>503</b>

---

# Chapter 1

## Introduction

### 1.1 Preliminaries

In recent years, a great deal of work has been done on various aspects of differential equations of third order. There are different problems concerning third-order differential equations which have drawn the attention of researchers throughout the world. Third-order differential equations describe many mathematical models of great interest in engineering, biology and physics. Equations of the form

$$x''' + a(t)x'' + b(t)x' + c(t)x = f(t)$$

arise in the study of entry-flow phenomenon, a problem of hydrodynamics which is of considerable importance in many branches of engineering (see [29]). Reynolds [51] has studied a model equation of the type

$$x'''(t) - \lambda x'(t) - 2x(t)x'(t) = \mu_1(Dx')(t) + \mu_2 \sin t, \quad (1.1a)$$

$$x(t + 2\pi) = x(t), \quad \int_{-\pi}^{\pi} x(t) dt = 0 \quad (1.1b)$$

which describes the steady flow of water in a long rectangular tank, oscillating horizontally near a resonant frequency. The integral damping operator  $D$  in (1.1a) is formally defined by

$$(Du)(t) = \left(\frac{2}{\pi}\right)^{1/2} \int_0^{\infty} u(t-s)s^{-1/2} ds$$

for all functions  $u$  with period  $2\pi$  and mean zero. Cox and Mortell [8] have considered (1.1a) and (1.1b) with  $(Dx')(t) = x(t)$ . In [28], Jackiewicz et al. have investigated the asymptotic behaviour of the solutions of Volterra integro-differential equations of the form

$$x'(t) = \gamma x(t) + \int_0^1 (\lambda + \mu t + \nu s)x(s) ds, \quad t \geq 0,$$
$$x(0) = 1$$

with the help of third-order differential equations of the type

$$x''' = \gamma x'' + (\lambda + (\mu + \nu)t)x' + (2\mu + \nu)x,$$

where  $\lambda$ ,  $\gamma$ ,  $\mu$  and  $\nu$  are real parameters and  $\nu + \mu \neq 0$ . In the development of a mathematical theory of thyroid-pituitary interaction, Danziger and Elemergreen (see [18], p. 133) have obtained the following third-order linear differential equations:

$$\alpha_3\theta''' + \alpha_2\theta'' + \alpha_1\theta' + (1+k)\theta = kc, \quad \theta < c$$

and

$$\alpha_3\theta''' + \alpha_2\theta'' + \alpha_1\theta' + \theta = 0, \quad \theta > c.$$

These equations describe the variation of thyroid hormone with time. Here  $\theta = \theta(t)$  is the concentration of thyroid hormone at time  $t$  and  $\alpha_3, \alpha_2, \alpha_1, k$  and  $c$  are constants. For other application of third-order differential equations, one may refer to Chap. 4 in [21].

In early 1950s, Alan Llyod Hodgkin and Andrew Huxley developed a mathematical model for the propagation of electrical pulses in the nerve of a squid. The original model describes the ionic mechanisms underlying the initiation and propagation of action potentials in the squid giant axon. The Hodgkin–Huxley model is a set of nonlinear differential equations that approximates the electrical characteristics of excitable cells such as neurons and cardiac myocytes. The model has played a vital role in biophysics and neuronal modelling. Alan Llyod Hodgkin and Andrew Huxley were awarded a Nobel prize in 1963 for this work. A reduced version of the Hodgkin–Huxley model was proposed by Nagumo. He suggested a third-order differential equation of the form

$$x''' - cx'' + f'(x)x' - \frac{b}{c}x = 0$$

as a model exhibiting many of the features of the Hodgkin–Huxley equations, where the function  $f$  is an entire function. For more details of Nagumo's equations, we refer the reader to the paper by McKeen [44] who gave some of the background of these equations and summarised some of the numerical results of this model. For application in physics, Vreeke and Sandquist [62] proposed the following system of differential equations (which is equivalent to a system of third-order differential equations)

$$\begin{aligned} \frac{dx_1}{dt} &= x_1(\gamma_1(1-x_2) + \gamma_2(1-x_3)), \\ \frac{dx_2}{dt} &= \gamma_3(x_1 - x_2), \\ \frac{dx_3}{dt} &= \gamma_4(x_1 - x_3) \end{aligned}$$

to describe the two temperature feedback nuclear reactor problem, where  $x_1$  is normalised neutron density,  $x_2$  and  $x_3$  are normalised temperatures,  $x_2$  being associated with fuel and  $x_3$  with the moderator or coolant,  $\gamma_3$  and  $\gamma_4$  are positive heat transfer coefficients,  $\gamma_1$  and  $\gamma_2$  are normalised effective neutron lifetime parameters associated with the temperature feedbacks, and the expression  $\gamma_1(1 - x_2) + \gamma_2(1 - x_3)$  in the first equation is called the *reactivity* and is a measure of multiplication factor of the neutrons in the fusion reactor.

The Kuramoto–Sivashinsky equation

$$u_t + u_{xxxx} + u_{xx} + \frac{1}{2}u^2 = 0$$

arises in a wide variety fascinating physical phenomena. For instance, we recall that the Kuramoto–Sivashinsky equation is introduced to describe pattern formulation in reaction diffusion systems, and to model the instability of flame front propagation (see Y. Kuramoto and T. Yamada [37] and D. Michelson [45]). To find the travelling wave solutions of this partial differential equation, we may use the substitution of the form  $u(x, ct) = u(x - ct)$  with period  $c$  and one has to solve the nonlinear third-order differential equation of the form

$$\lambda u'''(x) + u'(x) + f(u) = 0,$$

where  $\lambda$  is a parameter and  $f$  is an even function. On the other hand, by the qualitative behaviour of the above third-order differential equation, one can predict the behaviour of the above-mentioned partial differential equation. Hence, it would be interesting to study the qualitative behaviour of solutions of third-order linear and nonlinear differential equations. One may observe that the oscillation theory of second-order differential equations is well developed. It is only recently that the oscillation theory of third-order equations has been given a serious study.

In 1908, Birkhoff [6] applied a method of projective geometry and started the study of separation and comparison theorems for third-order equations. While Birkhoff's paper is a necessary reference to many papers on differential equations, his results or methods are seldom quoted. In 1961, Hanan [25] studied the oscillation and nonoscillation of two different types of third-order differential equation. The works due to Hanan [25] and Lazer [42] are the starting point of many investigations on the oscillation and asymptotic behaviour of solutions of third-order differential equations. In 1969, Barrett [5] made a self contained study on development of third-order equations. We may note that the third order is the lowest order where truly non-self-adjoint equations do occur. For example,  $x''' + tx' + t^2x = 0$  is a non-self-adjoint third-order differential equation. The self-adjoint form of the third-order equation is

$$x''' + q(t)x' + \frac{1}{2}q'(t)x = 0, \quad (1.2)$$

where  $q \in C^1([\sigma, \infty), R)$  and  $\sigma \in R$ . In the study of second- and fourth-order equations, the self-adjoint form of an equation plays a significant role. However, the

self-adjoint form of the third-order equation does not play any special role in the theory. The general solution of (1.2) is given by

$$x(t) = c_1 u^2(t) + c_2 u(t)v(t) + c_3 v^2(t),$$

where  $u(t)$  and  $v(t)$  are linearly independent solutions of the second-order equation

$$z'' + \frac{1}{4}q(t)z = 0,$$

and  $c_1, c_2$  and  $c_3$  are constants. Obviously  $u^2(t), u(t)v(t)$  and  $v^2(t)$  are linearly independent solutions of (1.2).

During last three decades, a considerable amount of work has been done on oscillation theory of linear homogeneous third-order differential equations with variable coefficients of the form

$$x''' + a(t)x'' + b(t)x' + c(t)x = 0, \quad (\text{LH1})$$

under various sign restrictions on the coefficient functions  $a(t), b(t)$  and  $c(t)$ , where  $a, b$  and  $c \in C([\sigma, \infty), \mathbb{R})$ . If  $a$  is twice continuously differentiable, then the transformation

$$x = y \exp\left(-\frac{1}{3} \int_{\sigma}^t a(s) ds\right)$$

transforms (LH1) to an equation of the form

$$y''' + Q(t)y' + P(t)y = 0, \quad (1.3)$$

where

$$P(t) = \frac{1}{27}[2a^3(t) - 9a(t)b(t) + 27c(t) - 9a''(t)]$$

and

$$Q(t) = \frac{1}{3}[3b(t) - 3a'(t) - a^2(t)].$$

Third-order differential equations of the form

$$(r(t)x'')' + q(t)x' + p(t)x = 0, \quad (\text{LH2})$$

where  $p, q, r \in C([\sigma, \infty), \mathbb{R})$  such that  $r(t) > 0$ , are more general than (LH1). In fact, (LH1) may be written as (LH2), where  $r(t) = \exp(\int_{\sigma}^t a(s) ds)$ ,  $q(t) = r(t)b(t)$  and  $p(t) = r(t)c(t)$ .

In [21], Greguš discussed oscillatory and asymptotic behaviour of solutions of equations of the type

$$x''' + 2A(t)x' + (A'(t) + B(t))x = 0. \quad (1.4)$$



Equation (1.4) is obtained from (1.3) by putting  $A(t) = \frac{1}{2}Q(t)$  and  $B(t) = P(t) - A'(t)$ . Greguš has assumed that  $a \in C^2([\sigma, \infty), R)$  and  $b \in C^1([\sigma, \infty), R)$ . The form (1.4) of a differential equation is called the *normal form* and the function  $B(t)$  is referred to as the *Laguerre invariant*. Although many authors (see [4, 11, 13, 14, 17, 19, 20, 25–27, 30–33, 36, 42, 55–57, 59]) have considered Eq. (1.3), it is always interesting to obtain sufficient conditions for oscillation/nonoscillation of solutions of (LH1) explicitly in terms of the coefficient functions  $a(t)$ ,  $b(t)$  and  $c(t)$ . For this reason, Greguš has discussed Eq. (LH1) in Chap. 2 in his monograph (see [21]). However, very little work has been done (see [9, 10, 47–49]) on the linear nonhomogeneous differential equations of third order of the type

$$x''' + a(t)x'' + b(t)x' + c(t)x = f(t), \quad (\text{NH1})$$

or

$$(r(t)x'')' + q(t)x' + p(t)x = f(t), \quad (\text{NH2})$$

where  $a, b, c, p, q, r \in C([\sigma, \infty), R)$  and  $r(t) > 0$ . Unlike second-order equations, (LH2) cannot be transformed to an equation of the form (1.3), and (NH1) cannot be transformed to an equation of the type

$$x''' + Q(t)x' + P(t)x = f(t).$$

Thus, the oscillation theory for Eqs. (LH2) and (NH2) is to be developed systematically.

A function  $x \in C([\sigma, \infty), R)$  is said to be nonoscillatory, if there exists a  $t_1 \geq \sigma$  such that  $x(t) > 0$  or  $< 0$  for  $t \geq t_1$ ; otherwise,  $x(t)$  is called oscillatory. By a solution of any of the above-mentioned differential equations, we mean a real-valued solution of that equation which exists on  $[T, \infty)$  and is nontrivial in any neighbourhood of infinity, where  $T \geq \sigma$  depends on the solution. Equation (LH1) is said to be nonoscillatory if all its solutions are nonoscillatory. It is said to be oscillatory if it admits an oscillatory solution. Equation (LH1) is said to be strongly oscillatory if all of its solutions are oscillatory. These definitions also hold good for other differential equations without delay. Such a classification of definitions is necessary because we have third-order differential equations which admit both oscillatory and nonoscillatory solutions. For example,  $x''' - x = 0$  admits solutions  $x_1(t) = e^t$  and  $x_2(t) = e^{-t/2} \sin \frac{\sqrt{3}}{2}t, t \geq 0$ . It is easy to see that all solutions of  $x''' - x' = 0$  are nonoscillatory. The nonhomogeneous equation

$$x''' - 2x' + e^t x = 1 + e^{-t}, \quad t \geq 0$$

has a nonoscillatory solution  $x(t) = e^{-t}$ . From Lazer's work (Theorem 1.3 in [42]), it follows that  $x''' - 2x' + e^t x = 0$  is oscillatory. Consequently, the above nonhomogeneous equation is oscillatory (see Corollary 4.2.4 in [9]). The equation

$$x''' - x = -\frac{9}{16}e^{-t/2}, \quad t \geq 0$$

admits a nonoscillatory solution,  $x_1(t) = \frac{1}{2}e^{-t/2}$ . Clearly,  $x_2(t) = e^{-t/2}(\frac{1}{2} + \sin \frac{\sqrt{3}}{2}t)$  is an oscillatory solution of the equation. Further, the nonhomogeneous delay differential equation

$$x''' - e^{\pi/2}x\left(t - \frac{\pi}{2}\right) = (e^{\pi/2} - 1)\cos t, \quad t \geq 0$$

admits an oscillatory solution  $x_1(t) = \sin t$  and a nonoscillatory solution  $x_2(t) = \sin t + e^t$ . The homogeneous delay differential equation

$$x'''(t) + 2x'(t) - x\left(t - \frac{3\pi}{2}\right) = 0$$

has an oscillatory solution  $x_1(t) = \sin t$ . If  $\phi(m) = m^3 + 2m - e^{-(3\pi/2)m} = 0$ , then  $\phi(0) = -1 < 0$  and  $\phi(m) \rightarrow \infty$  as  $m \rightarrow \infty$ . Hence there exists  $m_1 > 0$  such that  $\phi(m_1) = 0$ . Consequently,  $x_2(t) = e^{m_1 t}$  is a nonoscillatory solution of the equation.

It would be interesting to obtain an example of third-order differential equation which has only oscillatory solutions. Equation (LH1) is said to be disconjugate in  $[T, \infty)$ ,  $T \geq \sigma$  if no solution of (LH1) has more than two zeros in  $[T, \infty)$ , counting multiplicities. However, it carries no meaning for nonhomogeneous and nonlinear equations.

Now, we state oscillation of a solution of (LH1) in two different ways and show that these definitions are equivalent. Both definitions are used freely for linear homogeneous equations of the form (LH1) or (LH2). However, these definitions differ in case of linear nonhomogeneous and nonlinear equations.

**Definition 1.1.1** A solution  $x(t)$  of (LH1) is said to be oscillatory on  $[T_x, \infty)$ ,  $T_x \geq \sigma$ , if it has arbitrarily large zeros in  $[T_x, \infty)$ , that is, there exists a sequence  $\{t_n\} \subset [T_x, \infty)$  such that  $t_n \rightarrow \infty$  as  $n \rightarrow \infty$  and  $x(t_n) = 0$  for  $n = 1, 2, 3, \dots$

**Definition 1.1.2** A solution  $x(t)$  of (LH1) is said to be oscillatory on  $[T_x, \infty)$ ,  $T_x \geq \sigma$  if it has an infinite number of zeros in  $[T_x, \infty)$ .

These two definitions are equivalent. Suppose that a solution  $x(t)$  of (LH1) is oscillatory on  $[T_x, \infty)$  in the sense of Definition 1.1.2, then  $x(t)$  has infinite number of zeros in  $[T_x, \infty)$ . We claim that the infinite set  $A \subset [T_x, \infty)$  of zeros of  $x(t)$  is unbounded. If not, then  $A$  is bounded and hence there exist real numbers  $a$  and  $b$  ( $T_x \leq a < b < \infty$ ) such that  $A \subset [a, b] \subset [T_x, \infty)$ . From the Weierstrass theorem, it follows that  $A$  has a limit point, say  $t_0$ . Hence there exists a sequence  $\{t_n\} \subset A$  such that  $t_n \rightarrow t_0$  as  $n \rightarrow \infty$ . Thus  $t_0 \in [a, b]$  and  $x(t_n) = 0$  for  $n = 1, 2, 3, \dots$ . Since  $x$  is continuous,  $x(t_0) = 0$ . If  $x'(t_0) \neq 0$ , then  $x'(t_0) > 0$  or  $< 0$ . We may assume, without any loss of generality, that  $x'(t_0) > 0$ . From the continuity of  $x'(t)$ , it follows that there exists a  $\delta > 0$  such that  $x'(t) > 0$  for  $t \in (t_0 - \delta, t_0 + \delta)$ . Hence  $x(t)$  is increasing in  $(t_0 - \delta, t_0 + \delta)$ . Since  $x(t_0) = 0$ , we have  $x(t) < 0$  for  $t \in (t_0 - \delta, t_0)$  and  $x(t) > 0$  for  $t \in (t_0, t_0 + \delta)$ . On the other hand, there exists a  $t^* \in A$ ,  $t^* \neq t_0$  such

that  $t^* \in (t_0 - \delta, t_0 + \delta)$  as  $t_0$  is a limit point of  $A$ . Hence  $x(t^*) = 0$ , a contradiction. Thus  $x'(t_0) = 0$ . Further, if  $x''(t_0) \neq 0$ , then we may assume, without any loss of generality, that  $x''(t_0) > 0$ . Since  $x''(t)$  is continuous, there exists an  $\eta > 0$  such that  $x''(t) > 0$  for  $t \in (t_0 - \eta, t_0 + \eta)$ . Hence  $x'(t)$  is continuous and increasing in  $(t_0 - \eta, t_0 + \eta)$ . As  $x'(t_0) = 0$ , then  $x'(t) < 0$  for  $t \in (t_0 - \eta, t_0)$  and  $x'(t) > 0$  for  $t \in (t_0, t_0 + \eta)$ . Hence  $x(t)$  is decreasing in  $(t_0 - \eta, t_0)$  and increasing in  $(t_0, t_0 + \eta)$ . Since  $x(t_0) = 0$ , we have  $x(t) > 0$  for  $t \in (t_0 - \eta, t_0) \cup (t_0, t_0 + \eta)$ . However, there exists a  $t^{**} \neq t_0$ ,  $t^{**} \in A$  such that  $t^{**} \in (t_0 - \eta, t_0 + \eta)$  because  $t_0$  is a limit point of  $A$ . Consequently,  $0 < x(t^{**}) = 0$ , a contradiction. Hence  $x''(t_0) = 0$ . Thus from uniqueness it follows that  $x(t) \equiv 0$  on  $[T_x, \infty)$ , a contradiction. Then our claim holds, that is,  $A$  is unbounded. Hence there exists a sequence  $\{t_n\} \subset A$  such that  $t_n \rightarrow \infty$  as  $n \rightarrow \infty$ . As  $x(t_n) = 0$  for  $n = 1, 2, 3, \dots$ , then  $x(t)$  is oscillatory in the sense of Definition 1.1.1.

Conversely, if  $x(t)$  is oscillatory in the sense of Definition 1.1.1, then obviously, it is oscillatory in the sense of Definition 1.1.2.

These two definitions are equivalent for solutions of linear homogeneous second-order differential equations of the form

$$(r(t)x')' + p(t)x = 0,$$

but not equivalent for a function  $x \in C([\sigma, \infty), R)$ . We may note that the equation

$$x'' = t^2(30t^2 - 1) \sin \frac{1}{t} - 10t^3 \cos \frac{1}{t} + 21t^5, \quad t > 0$$

admits a solution  $x(t) = t^6(\sin \frac{1}{t} + \frac{t}{2})$ , which has an infinite number of zeros in  $(0, 1]$ , a finite number of zeros in  $[1, 2]$  and it is positive on  $(2, \infty)$ . Thus  $x(t)$  is oscillatory on  $(0, \infty)$  in the sense of Definition 1.1.2, but is nonoscillatory by Definition 1.1.1. This is because the equation is nonhomogeneous one. Further, one may observe that the zeros of an oscillatory solution of (LH1) are isolated. Indeed, if  $x(t)$  is an oscillatory solution of (LH1) on  $[T_x, \infty)$ , then there exists a sequence  $\{t_n\} \subset [T_x, \infty)$  such that  $t_n \rightarrow \infty$  as  $n \rightarrow \infty$  and  $x(t_n) = 0$ ,  $n = 1, 2, 3, \dots$ . We claim that  $t_n$  is isolated for all  $n$ . If not, then  $t_m$  is not isolated for some  $m \geq 1$ . Thus  $t_m$  is a limit point of  $\{t_n; n \geq 1\}$ . Hence there exists a subsequence  $\{t_{n_j}\}$  of  $\{t_n\}$  such that  $t_{n_j} \rightarrow t_m$  as  $j \rightarrow \infty$ . But  $t_n \rightarrow \infty$  as  $n \rightarrow \infty$  implies that  $t_{n_j} \rightarrow \infty$  as  $j \rightarrow \infty$ , a contradiction. Hence our assertion stands.

The adjoint of (LH1) is given by

$$((z' - a(t)z)' + b(t)z)' - c(t)z = 0. \quad (\text{LH1}^*)$$

Definitions 1.1.1 and 1.1.2 are equivalent for (LH1) {(LH1\*)}. However, if we consider (LH1) {(LH1\*)} on  $(0, d)$ , where  $d < \infty$ , then Definition 1.1.1 has no meaning. Here we repeat that a nontrivial solution  $x(t)$  of (LH1) {(LH1\*)} is said to be nonoscillatory if it is not oscillatory. If Eq. (LH1) {(LH1\*)} has a nontrivial oscillatory solution, then it is said to be oscillatory; otherwise, Eq. (LH1) {(LH1\*)} is said to be nonoscillatory.

Let  $S$  and  $S^*$  denote the solution space of (LH1) and (LH1\*), respectively. Thus each of them is a three-dimensional vector space over the field of real numbers. Let  $S_1 \{S_1^*\}$  denote a nontrivial subspace of  $S \{S^*\}$ . Then  $S_1 \{S_1^*\}$  is said to be nonoscillatory, if every nonzero member of  $S_1 \{S_1^*\}$  is nonoscillatory.  $S_1 \{S_1^*\}$  is said to be weakly oscillatory, if it contains a nontrivial oscillatory and nonoscillatory solution.  $S_1 \{S_1^*\}$  is said to be strongly oscillatory, if every nonzero member of  $S_1 \{S_1^*\}$  oscillates, and  $S_1 \{S_1^*\}$  is said to be oscillatory if  $S_1 \{S_1^*\}$  is either weakly oscillatory or strongly oscillatory. It may be noted that weakly oscillatory definition applies only to subspaces of dimension greater than or equal to two. If  $S \{S^*\}$  is nonoscillatory, weakly oscillatory or strongly oscillatory, then (LH1) {(LH1\*)} is said to be nonoscillatory, weakly oscillatory or strongly oscillatory, respectively.

## 1.2 Third-Order Linear Differential Equations with Constant Coefficients

Consider the linear third-order differential equation with constant coefficients of the form

$$x''' + ax'' + bx' + cx = 0, \quad t \geq \sigma, \quad (1.5)$$

where  $a, b$  and  $c$  are real constants. The auxiliary or the characteristic equation of (1.5) is given by

$$m^3 + am^2 + bm + c = 0. \quad (1.6)$$

The transformation  $n = m + \frac{a}{3}$  transforms (1.6) to

$$n^3 + 3Hn + G = 0, \quad (1.7)$$

where  $H = \frac{1}{3}(b - \frac{a^2}{3})$  and  $G = c - \frac{ab}{3} + \frac{2a^3}{27}$ .

From "Theory of Equations" (see [7]), it is well known that  $G^2 + 4H^3 > 0$  implies that (1.7) has two imaginary roots and a real root. If  $G^2 + 4H^3 \leq 0$ , then all the three roots of (1.7) are real. In particular,  $G^2 + 4H^3 = 0$  implies that two of these three roots are equal. If  $G = 0$  and  $H = 0$ , then all the three roots are equal. Two imaginary roots  $\alpha + i\beta$  and  $\alpha - i\beta$  of (1.7) give rise to two oscillatory solutions

$$x_1(t) = e^{(\alpha - \frac{a}{3})t} \cos \beta t \quad \text{and} \quad x_2(t) = e^{(\alpha - \frac{a}{3})t} \sin \beta t$$

of (1.5). It is well known that an algebraic equation of an odd degree has at least one real root of sign opposite to that of its last term. Consequently, Eq. (1.5) always admits a nonoscillatory solution of the type  $x_3(t) = e^{mt}$ , where  $m < 0$  if  $c > 0$  and  $m > 0$  if  $c < 0$ . If  $c = 0$ , then  $m = 0$ . Thus there is no linear homogeneous third-order differential equation with constant coefficients, which is strongly oscillatory.

According to the signs of  $a, b$  and  $c$ , one may consider the following eight different cases, viz., (i)  $a \geq 0, b \leq 0, c > 0$ , (ii)  $a \leq 0, b \leq 0, c > 0$ , (iii)  $a \leq 0, b \leq 0, c < 0$ , (iv)  $a \geq 0, b \leq 0, c < 0$ , (v)  $a \geq 0, b \geq 0, c > 0$ , (vi)  $a \leq 0, b \geq 0, c > 0$ , (vii)  $a \geq 0, b \geq 0, c < 0$  and (viii)  $a \leq 0, b \geq 0, c < 0$ .

**Proposition 1.2.1** *Suppose that  $a \geq 0$ ,  $b \leq 0$  and  $c > 0$ . Then the following holds:*

(i) *Equation (1.5) admits oscillatory solutions if and only if*

$$\frac{2a^3}{27} - \frac{ab}{3} + c - \frac{2}{3\sqrt{3}} \left( \frac{a^2}{3} - b \right)^{3/2} > 0 \quad (1.8)$$

*holds, and*

(ii) *When (1.8) is satisfied, all solutions of (1.5) are oscillatory except constant multiples of one solution which does not vanish on  $[\sigma, \infty)$  and which together with all of its derivatives is monotonic on  $[\sigma, \infty)$  and approaches zero as  $t$  tends to  $\infty$ . In fact, when (1.8) is satisfied, nonoscillatory solutions of (1.5) form a one-dimensional subspace of the solution space of (1.5).*

*Proof* (i) Clearly  $G > 0$  and  $H \leq 0$ . Further,  $G^2 + 4H^3 > 0$  if and only if  $(G - 2(-H)^{3/2})(G + 2(-H)^{3/2}) > 0$  if and only if  $G - 2(-H)^{3/2} > 0$ , that is (1.8) holds. Hence (1.8) holds if and only if (1.5) admits oscillatory solutions.

(ii) Since  $G > 0$  and (1.8) holds, (1.7) admits a negative root  $\gamma$  and two imaginary roots  $\alpha + i\beta$  and  $\alpha - i\beta$ . From ‘‘Theory of equations’’, it is clear that  $\gamma + (\alpha + i\beta) + (\alpha - i\beta) = 0$  holds and hence  $\gamma = -2\alpha$ . This in turn implies that  $\alpha > 0$ . Thus,

$$\left\{ e^{(\alpha - \frac{\gamma}{3})t} \cos \beta t, e^{(\alpha - \frac{\gamma}{3})t} \sin \beta t, e^{(\gamma - \frac{\gamma}{3})t} \right\}$$

is a basis of the solution space of (1.5). Clearly, constant multiples of  $e^{(\gamma - \frac{\gamma}{3})t}$  are nonoscillatory solutions of (1.5). These solutions and their derivatives are monotonic and tend to zero as  $t$  tends to  $\infty$ . To complete the proof of the proposition, it is enough to show that

$$x(t) = e^{(\alpha - \frac{\gamma}{3})t} (\lambda_1 \cos \beta t + \lambda_2 \sin \beta t) + \lambda_3 e^{(\gamma - \frac{\gamma}{3})t}$$

is oscillatory, where  $\lambda_1, \lambda_2$  and  $\lambda_3$  are constants such that  $\lambda_1$  and  $\lambda_2$  are not simultaneously equal to zero. If  $\lambda_3 = 0$ , then  $x(t)$  is oscillatory. Without any loss of generality, we assume that  $\lambda_3 > 0$ . If possible, let  $x(t) < 0$  for  $t > t_0$ . Since  $u(t) = \lambda_1 \cos \beta t + \lambda_2 \sin \beta t$  is oscillatory, there exists a  $t_1 > t_0$  such that  $u(t_1) = 0$ . So  $x(t_1) > 0$ , a contradiction. Suppose that  $x(t) > 0$  for  $t > t_0$ . So  $e^{(\alpha - \gamma)t} u(t) > -\lambda_3$  for  $t > t_0$ , a contradiction, because  $\alpha - \gamma > 0$  implies that  $e^{(\alpha - \gamma)t} u(t)$  oscillates between  $-\infty$  and  $\infty$ . This completes the proof.  $\square$

**Proposition 1.2.2** *Suppose that  $a \leq 0$ ,  $b \leq 0$  and  $c > 0$ . Then*

(i) *Equation (1.5) admits oscillatory solutions if and only if (1.8) is true, and*

(ii) *Further, if (1.8) is satisfied, then the conclusion of Proposition 1.2.1 remains true.*

*Proof* (i) Clearly,  $H \leq 0$ . (1.8) implies that  $G > 0$ ,  $G - 2(-H)^{3/2} > 0$  and hence  $G^2 + 4H^3 > 0$ . Consequently, (1.5) admits oscillatory solutions. On the other hand, if (1.5) admits oscillatory solutions, then (1.6) has two imaginary roots and a real

root. So (1.7) has two imaginary roots and a real root. This in turn implies that  $G^2 + 4H^3 > 0$ . Since  $c > 0$ , the real root of (1.6) is negative. Thus  $\gamma - \frac{a}{3} < 0$ . Since  $a < 0$ , the real root  $\gamma$  of (1.7) is negative. Consequently,  $G > 0$  and (1.8) holds.

(ii) If  $\alpha + i\beta$  and  $\alpha - i\beta$  are imaginary roots of (1.7), then  $\gamma + (\alpha + i\beta) + (\alpha - i\beta) = 0$  implies that  $\alpha > 0$  and

$$\{e^{(\alpha-\frac{a}{3})t} \cos \beta t, e^{(\alpha-\frac{a}{3})t} \sin \beta t, e^{(\gamma-\frac{a}{3})t}\}$$

defines a basis of the solution space of (1.5). The rest of the proof is same as that of Proposition 1.2.1.  $\square$

**Proposition 1.2.3** *Suppose that  $a \leq 0$ ,  $b \leq 0$  and  $c < 0$ . Then*

(i) *Equation (1.5) admits oscillatory solutions if and only if*

$$-\frac{2a^3}{27} + \frac{ab}{3} - c - \frac{2}{3\sqrt{3}} \left( \frac{a^2}{3} - b \right)^{3/2} > 0 \quad (1.9)$$

*holds,*

- (ii) *When (1.9) holds, oscillatory solutions of (1.5) form a two-dimensional subspace of the solutions space of (1.5), the zeros of any two oscillatory solutions of (1.5) separate on  $[\sigma, \infty)$  and the absolute values of the successive maxima and minima form a decreasing sequence, these oscillatory solutions tend to zero as  $t \rightarrow \infty$ , and*
- (iii) *Equation (1.5) admits a positive solution which tends to infinity as  $t$  tends to  $\infty$  and whose successive derivatives are positive.*

*Proof* (i) In this case  $G < 0$  and  $H \leq 0$ . Clearly, (1.9) holds if and only if (1.5) admits oscillatory solutions.

(ii) If (1.9) holds, then (1.7) has two imaginary roots, say  $\alpha + i\beta$  and  $\alpha - i\beta$  and a positive root  $\gamma$ , say. So  $\gamma + (\alpha + i\beta) + (\alpha - i\beta) = 0$  implies that  $\alpha < 0$  and

$$\{e^{(\alpha-\frac{a}{3})t} \cos \beta t, e^{(\alpha-\frac{a}{3})t} \sin \beta t, e^{(\gamma-\frac{a}{3})t}\}$$

defines a basis of the solution space of (1.5). The general solution of (1.5) is given by

$$x(t) = \lambda_1 e^{(\alpha-\frac{a}{3})t} \cos \beta t + \lambda_2 e^{(\alpha-\frac{a}{3})t} \sin \beta t + \lambda_3 e^{(\gamma-\frac{a}{3})t},$$

where  $\lambda_1, \lambda_2$  and  $\lambda_3$  are reals. We may note that  $x(t)$  is oscillatory, if and only if  $\lambda_3 = 0$ . Indeed,  $\lambda_3 = 0$  implies that  $x(t)$  is oscillatory. If  $\lambda_3 \neq 0$ , then

$$|x(t)| \geq |\lambda_3| e^{(\gamma-\frac{a}{3})t} - e^{(\alpha-\frac{a}{3})t} (|\lambda_1| + |\lambda_2|)$$

implies that  $x(t)$  is nonoscillatory. So oscillatory solutions of (1.5) form a two-dimensional subspace of the solution space of (1.5) with the basis

$$\{e^{(\alpha-\frac{a}{3})t} \cos \beta t, e^{(\alpha-\frac{a}{3})t} \sin \beta t\}.$$

Since the elements of the above basis are solutions of the second-order linear homogeneous equation

$$z'' - 2\left(\alpha - \frac{a}{3}\right)z' + \left[\left(\alpha - \frac{a}{3}\right)^2 + \beta^2\right]z = 0, \quad (1.10)$$

from Sturm's separation theorem, it follows that the zeros of any two oscillatory solutions of (1.5) separate on  $[\sigma, \infty)$ . Further,

$$2\left(\alpha - \frac{a}{3}\right)\left(\gamma - \frac{a}{3}\right) + \left(\alpha - \frac{a}{3}\right)^2 + \beta^2 = b$$

and  $b \leq 0$  implies that  $(\alpha - \frac{a}{3}) < 0$ . This, in turn, implies that the absolute values of the successive maxima and minima of an oscillatory solution of (1.5) form a decreasing sequence, and these oscillatory solutions tend to zero as  $t \rightarrow \infty$ .

(iii) Clearly,  $e^{(\gamma - \frac{a}{3})t}$  is the required nonoscillatory solution of (1.5) which together with all its derivatives tends to  $\infty$  as  $t \rightarrow \infty$ . This completes the proof of the proposition.  $\square$

**Proposition 1.2.4** *Suppose that  $a \geq 0$ ,  $b \leq 0$  and  $c < 0$ . Then the following hold:*

- (i) *Equation (1.5) admits oscillatory solutions, if and only if (1.9) holds,*
- (ii) *if (1.9) holds, then oscillatory solutions of (1.5) form a two-dimensional subspace of the solution space of (1.5). The zeros of any two linearly independent oscillatory solutions of (1.5) separate each other on  $[\sigma, \infty)$ , and these oscillatory solutions tend to zero as  $t \rightarrow \infty$ ,*
- (iii) *Equation (1.5) admits an oscillatory solution, if and only if all nonoscillatory solutions of (1.5) tend to  $\pm\infty$  as  $t \rightarrow \infty$ , and*
- (iv) *Equation (1.5) admits a positive solution, which tends to  $\infty$  as  $t \rightarrow \infty$  and whose successive derivative are positive and tend to  $\infty$  as  $t \rightarrow \infty$ .*

*Proof* (i) We may notice that  $H \leq 0$  and  $-G - 2(-H)^{3/2} > 0$  if and only if (1.9) holds. Clearly, (1.9) implies that  $G < 0$ . Since  $G^2 + 4H^3 = (-G - 2(-H)^{3/2})(-G + 2(-H)^{3/2})$ , (1.9) implies that  $G^2 + 4H^3 > 0$ . Thus, (1.7) has two imaginary roots and a real root. Consequently, (1.6) has two imaginary roots, say  $(\alpha - \frac{a}{3}) + i\beta$  and  $(\alpha - \frac{a}{3}) - i\beta$ , and a real root  $(\gamma - \frac{a}{3})$ , where  $\alpha + i\beta$ ,  $\alpha - i\beta$  and  $\gamma$  are the roots of (1.7). Thus (1.5) admits oscillatory solutions. On the other hand, if (1.5) admits an oscillatory solution, then (1.6) has two imaginary roots and a real root. This real root is positive because  $c < 0$ . Consequently,  $\gamma - \frac{a}{3} > 0$  because  $c < 0$ . Thus (1.7) has two imaginary roots and a positive root. Hence  $G^2 + 4H^3 > 0$  and  $G < 0$  because  $\gamma - \frac{a}{3} > 0$ . This in turn implies that (1.9) holds. Thus Eq. (1.5) admits an oscillatory solution, if and only if (1.9) holds.

(ii) If (1.9) holds, then a basis of the solution space of (1.5) is given by

$$\{e^{(\alpha - \frac{a}{3})t} \cos \beta t, e^{(\alpha - \frac{a}{3})t} \sin \beta t, e^{(\gamma - \frac{a}{3})t}\}.$$

If we set

$$x(t) = \lambda_1 e^{(\alpha - \frac{a}{3})t} \cos \beta t + \lambda_2 e^{(\alpha - \frac{a}{3})t} \sin \beta t + \lambda_3 e^{(\gamma - \frac{a}{3})t}, \quad (1.11)$$

where  $\lambda_1, \lambda_2$  and  $\lambda_3$  are reals such that  $\lambda_3 \neq 0$ , then  $x(t)$  is nonoscillatory because  $\gamma - \frac{a}{3} > 0$ . Again

$$\left(\alpha - \frac{a}{3}\right) + i\beta + \left(\alpha - \frac{a}{3}\right) - i\beta + \left(\gamma - \frac{a}{3}\right) = -a$$

implies that  $\alpha < 0$  and, we may write  $x(t)$  as

$$x(t) = e^{(\gamma - \frac{a}{3})t} [(\lambda_1 \cos \beta t + \lambda_2 \sin \beta t) e^{(\alpha - \gamma)t} + \lambda_3].$$

Hence, if (1.9) holds, then the oscillatory solutions of (1.5) form a two-dimensional subspace of the solution space of (1.5). Further, these oscillatory solutions of (1.5) tend to zero as  $t \rightarrow \infty$ , because  $\alpha - \frac{a}{3} < 0$ . Since  $e^{(\alpha - \frac{a}{3})t} \cos \beta t$  and  $e^{(\alpha - \frac{a}{3})t} \sin \beta t$  are solutions of (1.10), from Sturm's separation theorem, it follows that the zeros of any two linearly independent oscillatory solutions of (1.5) separate each other on  $[\sigma, \infty)$ .

(iii) If (1.5) admits an oscillatory solution and if  $x(t)$  is a nonoscillatory solution of (1.5), then we may write

$$x(t) = e^{(\alpha - \frac{a}{3})t} (\mu_1 \cos \beta t + \mu_2 \sin \beta t) + \mu_3 e^{(\gamma - \frac{a}{3})t},$$

where  $\mu_1, \mu_2$  and  $\mu_3$  are reals such that  $\mu_3 \neq 0$ . As

$$|x(t)| \geq |\mu_3| e^{(\gamma - \frac{a}{3})t} - (|\mu_1| + |\mu_2|) e^{(\alpha - \frac{a}{3})t},$$

then  $|x(t)| \rightarrow \infty$  as  $t \rightarrow \infty$ . Conversely, if every nonoscillatory solution of (1.5) tend to  $\pm\infty$  as  $t \rightarrow \infty$ , then (1.5) admits an oscillatory solution. Indeed, if all solutions of (1.5) are nonoscillatory, then all roots of (1.6) are positive. Hence the sum of the product of these roots taken two at a time is positive. But from Eq. (1.6), it follows that this sum is equal to  $b \leq 0$ , a contradiction.

(iv) Clearly,  $x(t) = e^{m_3 t}$ ,  $m_3 > 0$  is a nonoscillatory solution of (1.5) which together with all its derivatives tends to  $\infty$  as  $t \rightarrow \infty$ . We note that  $m_3 > 0$ , because  $c < 0$ . Hence the proposition is proved.  $\square$

**Proposition 1.2.5** *Suppose that  $a \geq 0$ ,  $b \geq 0$  and  $c > 0$ . Then*

- (i) *Equation (1.5) admits a nonoscillatory solution which together with all of its derivatives tend to zero as  $t \rightarrow \infty$ ,*
- (ii) *if  $3b > a^2$ , then (1.5) is oscillatory,*
- (iii) *if  $3b \leq a^2$ , then (1.5) is oscillatory if and only if*

$$\left| \frac{2a^3}{27} - \frac{ab}{3} + c \right| - \frac{2}{3\sqrt{3}} \left( \frac{a^2}{3} - b \right)^{3/2} > 0 \quad (1.12)$$

*holds,*



- [\*The Three-Cornered World.pdf\*](#)
- [\*\*Sailing Made Easy.pdf\*\*](#)
- [Ideas About Art.pdf, azw \(kindle\), epub, doc, mobi](#)
- [La mala hora online](#)
- [download online Lore and Science in Ancient Pythagoreanism](#)
- [read online Naakte zielen \(Deleu, Book 1\)](#)
  
- <http://jaythebody.com/freebooks/The-Three-Cornered-World.pdf>
- <http://www.celebritychat.in/?ebooks/Numerical-Methods-with-Worked-Examples--Matlab-Edition--2nd-Edition-.pdf>
- <http://academialanguagebar.com/?ebooks/Ideas-About-Art.pdf>
- <http://berttrotman.com/library/60-Hikes-Within-60-Miles--Seattle.pdf>
- <http://cavalldecartro.highlandagency.es/library/The-Expeditions.pdf>
- <http://cambridgebrass.com/?freebooks/Dragon-s-Egg--Cheela--Book-1-.pdf>